

MATHEMATICS AND EDUCATION IN MATHEMATICS, 1986

Proceedings of 15th Spring Conference of
the Union of Bulgarian Mathematicians,
Sunny Beach, April 2 - 6, 1986

Multipliers of integrals of Cauchy - Stieltjes type

Peyo Stoilov

Abstract

Let $\mathbb{G}$ be a domain with closed rectifiable Jordan curve ℓ . Let $K(\mathbb{G})$ be the space of all analytic functions in $\mathbb{G}$ representable by a Cauchy - Stieltjes integral. Let $\mathfrak{M}(K)$ be the class of all multipliers of the space $K(\mathbb{G})$. In this paper we prove that if f is bounded analytic function on $\mathbb{G}$ and

$$\operatorname{ess\,sup}_{\eta \in \ell} \int_{\ell} \frac{|f(\zeta) - f(\eta)|}{|\zeta - \eta|} |d\zeta| < \infty,$$

then $f \in \mathfrak{M}(K)$.

If $\mathbb{G} = \mathbb{D}$ is the unit disc, this theorem was proved for the first time by V. P. Havin. In particular for a smooth curve ℓ we prove that if $f' \in E^p(\mathbb{G})$, $p > 1$, then $f \in \mathfrak{M}(K)$, where $E^p(\mathbb{G})$ are the spaces of Smirnov.

1 Introduction

Let ℓ be a closed rectifiable Jordan curve bounding the finite and infinite domains $\mathbb{G}$ and $\mathbb{G}^-$ in the complex plane $\mathbb{C}$. Let ℓ_r ($0 < r < 1$) be the map of circle $|z| = r$ under conformal mapping $\zeta = \varphi(z)$ of the unit disc $\mathbb{D}$ on the domain $\mathbb{G}$. Let denote by $A(\mathbb{Q})$ the set of all analytic functions in a domain $\mathbb{Q} \subset \mathbb{C}$. For $0 < p \leq \infty$, let $E^p(\mathbb{G})$ be the Smirnov class [1,2] of functions $f \in A(\mathbb{G})$, for which

$$\|f\|_{E^p}^p = \sup_{0 < r < 1} \frac{1}{2\pi} \int_{\ell_r} |f(\zeta)|^p |d\zeta| < \infty, \quad 0 < p < \infty,$$

⁰**1980 Mathematics Subject Classification:** Primary 30E20, 30D50.

⁰**Key words and phrases:** Cauchy - Stieltjes integral, Multipliers, Spaces of Smirnov, Theorem of Smirnov - Kotchine, Closed Jordan curve.

$$\|f\|_{E^\infty} = \sup_{\zeta \in \mathbb{G}} |f(\zeta)| < \infty, \quad p = \infty.$$

If $\mathbb{G} = \mathbb{D}$, $\mathbb{D} = \{\zeta : |\zeta| < 1\}$, then $E^p(\mathbb{G}) = H^p$ is the usual Hardy class.

Let N denote the Nevanlinna class:

$$f \in N \Leftrightarrow f \in A(\mathbb{D}) : \quad \lim_{r \rightarrow 1^-} \int_{|\zeta|=1} \log^+ |f(r\zeta)| |d\zeta| < \infty,$$

and N^+ - the Smirnov class:

$$f \in N^+ \Leftrightarrow f \in N : \quad \lim_{r \rightarrow 1^-} \int_{|\zeta|=1} \log^+ |f(r\zeta)| |d\zeta| = \int_{|\zeta|=1} \log^+ |f(\zeta)| |d\zeta|.$$

Some basic facts.

$$1. f(\zeta) \in E^p(\mathbb{G}) \Leftrightarrow f(\varphi(z)) \sqrt[p]{\varphi'(z)} \in H^p;$$

2. Theorem of Smirnov - Kotchine:

If $f \in N^+$ and has L^p boundary values, then $f \in H^p$.

Let $M(\ell)$ is the space of all finite, complex Borel measures on ℓ with the usual variation norm. We denote by $K(\mathbb{G})$ the space of all functions $g \in A(\mathbb{G})$, representable by an integral of Cauchy - Stieltjes type

$$g(z) = \int_{\ell} \frac{d\mu(\zeta)}{\zeta - z} \stackrel{def}{=} K_{\mu}(z), \quad \mu \in M(\ell).$$

If $g \in K(\mathbb{G})$, then M_g will denote the set of all measures $\mu \in M(\ell)$ participating in the representation of g .

Since $K_{\mu}(z) = 0, \forall z \in \mathbb{G} \Leftrightarrow K_{\mu}(z) \in E^1(\mathbb{G}^-)$ and

$$E^1(\mathbb{G}^-) = \left\{ K_{\mu} : d\mu = f(\zeta)d\zeta, f(\zeta) \in L^1(\ell), \int_{\ell} \zeta^n f(\zeta)d\zeta = 0, n = -1, -2, \dots \right\},$$

then the space $K(\mathbb{G})$ is the Banach space with the natural norm

$$\|g\|_{K(\mathbb{G})} = \inf \left\{ \|\mu\|_{M(\ell)} : \mu \in M_g \right\},$$

isometrically isomorphic to $M(\ell)/E^1(\mathbb{G}^-)$.

Let $C_A(\overline{\mathbb{G}^-})$ denote the space of all functions h , analytic in $\mathbb{G}^-$ and continuous on $\overline{\mathbb{G}^-} = \mathbb{G}^- \cup \ell$, with norm

$$\|h\|_\infty = \sup \{ |h(z)| : z \in \mathbb{G}^- \}.$$

Since $M(\ell)/E^1(\mathbb{G}^-) \cong C_A^*(\mathbb{G}^-)$, then $K(\mathbb{G}) \cong C_A^*(\mathbb{G}^-)$ and

$$\|g\|_{K(\mathbb{G})} = \sup \left\{ \left| \int_\ell h(\eta) d\mu(\eta) \right| : h \in C_A(\mathbb{G}^-), \|h\|_\infty \leq 1 \right\}, \quad \mu \in M_g.$$

Taking into account that the set $A_0 = A(\overline{\mathbb{G}^-})$ is dense in $C_A(\mathbb{G}^-)$, then easily we receive for $h \in A_0$, $g = K_\mu$:

$$\begin{aligned} \left| \int_{\ell_r} h(\zeta) g(\zeta) d\zeta \right| &= \left| \int_{\ell_r} h(\zeta) \left(\int_\ell \frac{d\mu(\eta)}{\eta - \zeta} \right) d\zeta \right| = \\ &= \left| \int_\ell \left(\int_{\ell_r} \frac{h(\zeta)}{\zeta - \eta} d\zeta \right) d\mu(\eta) \right| = \left| \int_\ell h(\eta) d\mu(\eta) \right| \end{aligned}$$

and consequently

$$\|g\|_{K(\mathbb{G})} = \sup \left\{ \lim_{r \rightarrow 1^-} \left| \int_{\ell_r} h(\zeta) g(\zeta) d\zeta \right| : h \in A_0, \|h\|_\infty \leq 1 \right\}.$$

Let $\mathfrak{M}(K)$ be the class to all multipliers of the space $K(\mathbb{G})$:

$$\mathfrak{M}(K) = \{ f \in A(\mathbb{G}) : fg \in K(\mathbb{G}), \quad \forall g \in K(\mathbb{G}) \}.$$

Lemma. *If $f \in \mathfrak{M}(K)$, then*

$$\sup \left\{ \|fg\|_{K(\mathbb{G})} : g \in K(\mathbb{G}), \|g\|_{K(\mathbb{G})} \leq 1 \right\} < \infty.$$

Proof. Let $f \in \mathfrak{M}(K)$. We will show that the operator $g \rightarrow fg$ is bounded in $K(\mathbb{G})$. By the closed graph theorem it suffices to show that the graph of the operator $g \rightarrow fg$ is closed on $K(\mathbb{G})$.

Suppose $(g_n)_{n \geq 0} \subset K(\mathbb{G})$ with $g_n \rightarrow g$ and $fg_n \rightarrow \phi$ in the $K(\mathbb{G})$ -norm.

If $F \in K(\mathbb{G})$, then for all $z \in \mathbb{G}$,

$$F(z) = \int_\ell \frac{d\mu(\zeta)}{\zeta - z} \Rightarrow |F(z)| \leq (\inf |\zeta - z| : \zeta \in \ell)^{-1} \|F\|_{K(\mathbb{G})}.$$

From this inequality follows that the converse in norm in $K(\mathbb{G})$ implies a pointwise converse on $\mathbb{G}$. Consequently $\phi = fg$ and the graph of the operator $g \rightarrow fg$ is closed on $K(\mathbb{G})$. $\square$

From proved lemma follows that $\mathfrak{M}(K)$ is commutative Banach algebra with norm defined by

$$\|f\|_{\mathfrak{M}(K)} = \sup \left\{ \|fg\|_{K(\mathbb{G})} : g \in K(\mathbb{G}), \|g\|_{K(\mathbb{G})} \leq 1 \right\}.$$

If $\mathbb{G} = \mathbb{D}$, then the study of the space $\mathfrak{M} = \mathfrak{M}(K)$ was started in papers of V. P. Havin [3, 4] and was continued in [5, 6, 7]. For example, in [5, 6] was proved:

If f is bounded analytic function on $\mathbb{D}$ and

$$\operatorname{ess\,sup}_{|\eta|=1} \int_{|\zeta|=1} \frac{|f(\zeta) - f(\eta)|}{|\zeta - \eta|} |d\zeta| < \infty,$$

then $f \in \mathfrak{M}$.

In this paper this result is generalized for the multipliers of the space $K(\mathbb{G})$.

2 Main results

Theorem 1. *Let $\mathbb{G}$ be a domain with closed rectifiable Jordan curve ℓ .*

If $f \in E^\infty(\mathbb{G})$ and

$$\Lambda(f) \stackrel{\text{def}}{=} \operatorname{ess\,sup}_{\eta \in \ell} \int_{\ell} \frac{|f(\zeta) - f(\eta)|}{|\zeta - \eta|} |d\zeta| < \infty,$$

then $f \in \mathfrak{M}(K)$ and

$$\|f\|_{\mathfrak{M}(K)} \leq \|f\|_{E^\infty} + \Lambda(f).$$

Proof. Let $f \in E^\infty(\mathbb{G})$ and $\Lambda(f) < \infty$. Let $\mathbb{E} \subseteq \ell$ be a subset with total Lebesgue measure ($m(\mathbb{E}) = m(\ell)$) lying on ℓ such that

$$\|f\|_{E^\infty} = \sup_{\eta \in \mathbb{E}} |f(\eta)|.$$

Then

$$\begin{aligned} \|f\|_{\mathfrak{M}(K)} &= \sup \left\{ \|fg\|_{K(\mathbb{G})} : g \in K(\mathbb{G}), \|g\|_{K(\mathbb{G})} \leq 1 \right\} = \\ &= \sup \left\{ \lim_{r \rightarrow 1^-} \left| \int_{\ell_r} f(\zeta) g(\zeta) h(\zeta) d\zeta \right| : g \in K(\mathbb{G}), \|g\|_{K(\mathbb{G})} \leq 1, h \in A_0, \|h\|_\infty \leq 1 \right\}, \end{aligned}$$

where ℓ_r ($0 < r < 1$) is the map of circle $|z| = r$ under conformal mapping $\zeta = \varphi(z)$ of the unit disc $\mathbb{D}$ on the domain $\mathbb{G}$.

If $g \in K(\mathbb{G})$, then

$$g(z) = \int_{\ell} \frac{d\mu(\eta)}{\eta - z}, z \in \mathbb{G}$$

and

$$\begin{aligned} \left| \int_{\ell_r} f(\zeta)g(\zeta)h(\zeta)d\zeta \right| &= \left| \int_{\ell_r} f(\zeta)h(\zeta) \left(\int_{\ell} \frac{d\mu(\eta)}{\eta - \zeta} \right) d\zeta \right| = \\ &= \left| \int_{\ell} \left(\int_{\ell_r} \frac{f(\zeta)h(\zeta)}{\zeta - \eta} d\zeta \right) d\mu(\eta) \right| \leq \\ &\leq \int_{\ell} |d\mu(\eta)| \sup \left\{ \left| \int_{\ell_r} \frac{f(\zeta)h(\zeta)}{\zeta - \eta} d\zeta \right| : \eta \in \mathbb{E} \right\}, \end{aligned}$$

$\Rightarrow$

$$\left| \int_{\ell_r} f(\zeta)g(\zeta)h(\zeta)d\zeta \right| \leq \|g\|_{K(\mathbb{G})} \sup \left\{ \left| \int_{\ell_r} \frac{f(\zeta)h(\zeta)}{\zeta - \eta} d\zeta \right| : \eta \in \mathbb{E} \right\}.$$

Then

$$\|f\|_{\mathfrak{M}(K)} \leq \sup \left\{ \lim_{r \rightarrow 1^-} \left| \int_{\ell_r} \frac{f(\zeta)h(\zeta)}{\zeta - \eta} d\zeta \right| : \eta \in \mathbb{E}, h \in A_0, \|h\|_{\infty} \leq 1 \right\}.$$

Further, if $\eta \in \mathbb{E}$, $h \in A_0$, $\|h\|_{\infty} \leq 1$, we have

$$\begin{aligned} \left| \int_{\ell_r} \frac{f(\zeta)h(\zeta)}{\zeta - \eta} d\zeta \right| &\leq \left| \int_{\ell_r} \frac{f(\zeta) - f(\eta)}{\zeta - \eta} h(\zeta) d\zeta \right| + |f(\eta)| \left| \int_{\ell_r} \frac{h(\zeta)}{\zeta - \eta} d\zeta \right| \leq \\ &\leq \|h\|_{L^{\infty}(\ell_r)} \int_{\ell_r} \left| \frac{f(\zeta) - f(\eta)}{\zeta - \eta} \right| |d\zeta| + \|f\|_{E^{\infty}}. \end{aligned}$$

Then

$$\|f\|_{\mathfrak{M}(K)} \leq \sup \left\{ \lim_{r \rightarrow 1^-} \int_{\ell_r} \left| \frac{f(\zeta) - f(\eta)}{\zeta - \eta} \right| |d\zeta| : \eta \in \mathbb{E} \right\} + \|f\|_{E^{\infty}}.$$

We denote for $\eta \in \mathbb{E}$

$$F_{\eta}(\zeta) = \frac{f(\zeta) - f(\eta)}{\zeta - \eta}, \quad \zeta \in \mathbb{G}.$$

Then

$$\|f\|_{\mathfrak{M}(K)} \leq \sup \left\{ \|F_{\eta}(\zeta)\|_{E^1} : \eta \in \mathbb{E} \right\} + \|f\|_{E^{\infty}}.$$

To end the proof it is necessary to show that

$$\Lambda(f) < \infty \Rightarrow \sup_{\eta \in \mathbb{E}} \|F_\eta\|_{E^1} < \infty.$$

Let $\zeta = \varphi(z)$ is conformal mapping of the unit disc $\mathbb{D}$ on the domain $\mathbb{G}$, $\eta \in \mathbb{E}$ and $\eta = \varphi(t)$, $|t| = 1$. Then

$$F_\eta(\zeta) = \frac{f(\zeta) - f(\eta)}{\zeta - \eta} \in E^1(\mathbb{G}) \Leftrightarrow \Omega(z) \stackrel{def}{=} \frac{f(\varphi(z)) - f(\varphi(t))}{\varphi(z) - \varphi(t)} \varphi'(z) \in H^1.$$

In [8] was proved that the functions

$$\frac{1}{\varphi(z) - \varphi(t)} \in N^+.$$

Since

$$f(\varphi(z)) - f(\varphi(t)) \in H^\infty, \varphi'(z) \in H^1, \text{ then } \Omega(z) \in N^+.$$

Besides

$$\int_{|z|=1} |\Omega(z)| |dz| = \int_{\ell} |F_\eta(\zeta)| |d\zeta| \leq \Lambda(f) < \infty$$

and according to the Theorem of Smirnov - Katchine

$$\Omega(z) \in H^1, \|\Omega(z)\|_{H^1} \leq \Lambda(f) < \infty.$$

Consequently

$$\|F_\eta\|_{E^1} = \|\Omega(z)\|_{H^1} \leq \Lambda(f) < \infty$$

and

$$\|f\|_{\mathfrak{M}(K)} \leq \Lambda(f) + \|f\|_{E^\infty} < \infty.$$

□

Theorem 2. Let $\mathbb{G}$ be a domain with a smooth curve ℓ .

If $f \in E^\infty(\mathbb{G})$ and $f' \in E^p(\mathbb{G})$, $p > 1$, then $f \in \mathfrak{M}(K)$.

Proof. Let $f' \in E^p(\mathbb{G})$, $p > 1$, $z = z(s)$ ($0 \leq s \leq s_0$) is the equation of ℓ , where the arc length is as parameter. We shall prove that

$$\Lambda(f) = \sup_{\sigma \in [0, s_0]} \int_0^{s_0} \frac{|f(s) - f(\sigma)|}{|z(s) - z(\sigma)|} ds < \infty,$$

where $f(s) = f \circ z(s)$.

Since ℓ is a smooth curve, then

$$|z(s) - z(\sigma)| \geq |s - \sigma|,$$

where c_0 is a constant. Then

$$\begin{aligned} \int_0^{s_0} \frac{|f(s) - f(\sigma)|}{|z(s) - z(\sigma)|} ds &\leq \frac{1}{c_0} \int_0^{s_0} \frac{1}{|s - \sigma|} \left| \int_\sigma^s f'(x) dx \right| ds \leq \\ &\leq \frac{1}{c_0} \int_0^{s_0} \left(\int_\sigma^s |f'(x)|^p dx \right)^{1/p} |s - \sigma|^{-1/p} ds \leq \text{Const.} \|f'\|_{E^p(\mathbb{G})}. \end{aligned}$$

Consequently

$$\Lambda(f) \leq \text{Const.} \|f'\|_{E^p(\mathbb{G})} < \infty$$

and by Theorem 1 $f \in \mathfrak{M}(K)$. $\square$

Remark. It should be noted that if $\mathbb{G} = \mathbb{D}$, Theorem 2 remains valid and for $p = 1$ (Theorem of S. A. Vinogradov [5]: $f'(z) \in H^1 \Rightarrow f \in \mathfrak{M}$).

Whether Theorem 2 for $p = 1$ is correct, if $\mathbb{G}$ is domain with a smooth boundary, remains unknown.

References

- [1] I. I. Privalov. *Boundary properties of analytic functions*. GATT, Moscow, 1950. (Russian) German transl., VEB Deutscher Verlag, Berlin, 1956. MR0047765 (13, 926h)
- [2] P. L. Duren. *Theory of H^p spaces*. Academic Press, New York, 1970. MR0268655 (42 #3552)
- [3] V. P. Havin. *On analytic functions representable by an integral of Cauchy-Stieltjes type*. Vestnik Leningrad. Univ. Ser. Mat. Meh. Astronom. 13 no.1, 1958, 66-79. (Russian) MR0095256 (20 #1762)
- [4] V. P. Havin. *Relations between certain classes of functions regular in the unit disk*. Vestnik Leningrad. Univ. Ser. Mat. Meh. Astronom. 17 no.1, 1962, 102-110. (Russian) MR0152660 (27 #2635)
- [5] S. A. Vinogradov. *Properties of multipliers of Cauchy - Stieltjes integrals and some factorization problems for analytic functions*. Amer. Math. Sos. Transl. (2) vol. 115, 1980, 1-32. MR0586560 (58 #28518)

- [6] S. A. Vinogradov, M. G. Goluzina, V. P. Havin. *Multipliers and divisors of Cauchy-Stieltjes type integrals*. Zap. Nauen. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) 19, 1970, 55-78, 1970. (Russian) MR0291471 (45 #562)
- [7] S. V. Hruščëv, S. A. Vinogradov. *Inner functions and multipliers of Cauchy type integrals*. Ark. mat, 19, 1981, 23-42. MR0625535 (83c:30027)
- [8] G. C. Tumarkin. *Properties of analytic functions representable by integrals of Cauchy-Stieltjes and Cauchy-Lebesgue type*. Izv. Akad. Nauk Armjan. SSR Ser. Fiz.- Mat. Nauk 16 no. 5, 1963, 23-45. (Russian) MR0156955 (28 #197)

Department of Mathematics
Technical University
25, Tsanko Dustabanov,
Plovdiv, Bulgaria
e-mail: peyyyo@mail.bg